

It appears that you are currently using **Ad Blocking** software. What are the consequences? [Click here to learn more.](#)

This website uses cookies to ensure you get the best experience on our website. [More info](#)

[Nanotechnology](#) [Physics](#) [Earth](#) [Astronomy & Space](#) [Technology](#) [Chemistry](#) [Biology](#) [Other Sciences](#)

[Home](#) [Biology](#) [Ecology](#) January 11, 2018

# Lethal management of wolves in one place may make things worse nearby

January 11, 2018 by Mason Muerhoff, University of Wisconsin-Madison



Killing wolves to prevent predation on livestock may protect one farm but harm neighbors, according to preliminary conclusions of UW-Madison researchers. Credit: National Park Service

Lethal management of wolves following wolf attacks on livestock may have unintended consequences, a new study by researchers at the University of Wisconsin-Madison suggests.

When a wolf was killed under government programs to address wolf predation near a farm that had recently reported an attack, the authors of the study say two things happened: That farm experienced a decrease in the likelihood of livestock losses to wolves, but surrounding farms showed an increase.

Adrian Treves and Francisco J. Santiago-Avila regard their conclusions, published Jan. 10, 2018 in the journal *PLOS ONE*, as preliminary given the nature of the experimental data, which they did not collect, and because nonlethal interventions served as a pseudo-control group for comparison. According to Santiago-Avila, the study is an attempt to apply a higher standard

Featured	Last comments	Popular
	Surprising result shocks scientists studying spin	Jan 08, 2018 26
	Scientists take viewers to the center of the Milky Way	Jan 10, 2018 0
	Chemists discover plausible recipe for early life on Earth	Jan 08, 2018 53
	Physicists create first direct images of the square of the wave function of a hydrogen molecule	Jan 09, 2018 11
	Device creates negative mass—and a novel way to generate lasers	Jan 10, 2018 22

[more »](#)

Phys.org

 [Follow](#)

Phys.org on facebook

[Follow](#) 1.3M people are following [Phys.org](#). [Sign Up](#) to see who your friends are following.

Email newsletter

of science to an age-old conflict between carnivores and farmers.

"We wanted to evaluate if lethal management in a particular area once there is an attack by wolves on livestock in that area is actually effective in mitigating that attack," says Santiago-Avila. "Or another alternative would be if this is actually making things worse."

Treves is a professor at UW–Madison's Nelson Institute for Environmental Studies and head of the Carnivore Coexistence Lab. Santiago-Avila, his graduate student, is lead author of the study.

Treves and Santiago-Avila analyzed data acquired by the Little River Band of Ottawa Indians in Michigan. The data was collected by the Michigan Department of Natural Resources and consisted of 230 cases of wolf attacks on livestock in the Upper Peninsula from 1998 to 2014. Of those cases, 199 resulted in some form of intervention from the DNR. Thirty-one of the interventions involved killing a wolf, while 168 employed nonlethal methods for deterrence. Nonlethal methods include giving the owner advice, using guard dogs, or using fladry—colored flags that hang from fences to deter wolves from crossing them.

Treves and Santiago-Avila used three spatial scales around the site of an attack, first at one square mile, then at 26 square miles, and up to 320 square miles, in order to detect any "spillover effects" after lethal intervention.

The results showed farms that had a wolf killed experienced a 27 percent decrease in risk of another attack, but this was offset by a 22 percent increase at a number of farms in the same township—making the net benefit zero, according to the study.

One factor that could explain this spillover effect, says Santiago-Avila, is that wolves have strictly ordered social lives. Because each animal has a different role in the pack, killing one wolf may cause the pack to split up and spread out into new territory, potentially seeking food that requires fewer members to hunt. But the effect over all of the years captured in the study remained constant, reducing the likelihood that distribution was a main factor.



#### Relevant PhysicsForums posts

Did Neanderthals have the intellectual capacity to read? 3 hours ago

Examination project of EEG, Opinions and feedback 5 hours ago

Worm Species without Males Loses 25% of Genes! 18 hours ago

Do most caucasian homo sapiens have some Neanderthal DNA? 19 hours ago

Bird ID, please (seen at at the Sweetwater Wetlands in Tucson) Jan 10, 2018

Mid-season additions to flu vaccinations? Jan 10, 2018

More from Biology and Medical

Other explanations may account for it, too, but in order to conduct an experiment that meets the gold standard of science and draws stronger conclusions, Santiago-Avila says, changes to how the data are reported and tracked are necessary.

"It starts with accurate and sophisticated monitoring of what you're doing on the ground," Santiago-Avila said. "And that's one of the main concerns that the Little River Band had, that they couldn't see how they would be evaluating the data or if there was an intention to do so."

### Explore further: [Wolves to the slaughter: France approves cull to save sheep](#)

Journal reference: [PLoS ONE](#)

0 shares

Provided by: [University of Wisconsin-Madison](#)

[feedback to editors](#)

45 Like Tweet 14 Favorites Email Print PDF

#### Related Stories



### Wolves to the slaughter: France approves cull to save sheep

July 20, 2017


The French government on Thursday gave the green light for the cull of dozens of wolves in mountainous areas where sheep are under sustained attack.

### Livestock risks from Wisconsin wolves localized, predictable

June 1, 2011

It's an issue that crops up wherever humans and big predators — wolves. bears. lions — coexist.


#### Recommended for you



### Scientists make cells that enable the sense of touch

January 11, 2018

Researchers at the Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research at UCLA have, for the first time, coaxed human stem cells to become sensory interneurons—the cells that give us our sense of ...



### Researchers map druggable genomic targets in evolving malaria parasite

January 11, 2018

Researchers at University of California San

#### User comments

Please sign in to add a comment. Registration is free, and takes less than a minute. [Read more](#)

[Sign in](#)

---

[Click here to reset your password.](#)

[Sign in to get notified via email when new comments are made.](#)

- 
- |                                |                         |                                   |                                 |                                   |                         |
|--------------------------------|-------------------------|-----------------------------------|---------------------------------|-----------------------------------|-------------------------|
| <a href="#">Top</a>            | <a href="#">Help</a>    | <a href="#">Science X Account</a> | <a href="#">Feature Stories</a> | <a href="#">Android app</a>       | <a href="#">Connect</a> |
| <a href="#">Home</a>           | <a href="#">FAQ</a>     | <a href="#">Sponsored Account</a> | <a href="#">Latest news</a>     | <a href="#">iOS app</a>           |                         |
| <a href="#">Search</a>         | <a href="#">About</a>   | <a href="#">Newsletter</a>        | <a href="#">Week's top</a>      | <a href="#">Amazon Kindle</a>     |                         |
| <a href="#">Mobile version</a> | <a href="#">Contact</a> | <a href="#">RSS feeds</a>         | <a href="#">Archive</a>         | <a href="#">Push notification</a> |                         |